
Honors PreCalculus

Section 9.5: Series

Give it a try:

Find the sum of the natural numbers
1 to 100. Without a calculator 😊

$$\begin{array}{r} 1 + 2 + 3 + \dots + 99 + 100 = ? \\ + 100 + 99 + 98 + \dots + 2 + 1 \\ \hline 101 \quad 101 \end{array}$$

$$(101)100 = 10,100/2 = 5050$$

5050

Summation Notation

- The sum of the terms of a sequence $a_k = \{a_1, a_2, a_3, \dots, a_n\}$ can be represented as

$$\sum_{k=1}^n a_k$$

“The sum of a_k from $k = 1$ to $k = n$ ”

Summation Notation

- Find the sums:

$$\sum_{k=1}^5 3k = 3(1) + 3(2) + 3(3) + 3(4) + 3(5) = 45$$

$$\sum_{k=5}^8 k^2 = 5^2 + 6^2 + 7^2 + 8^2 = 174$$

$$\sum_{n=0}^4 \cos n\pi = \overset{|}{\cos 0} + \overset{-}{\cos \pi} + \overset{|}{\cos 2\pi} + \overset{-}{\cos 3\pi} + \overset{|}{\cos 4\pi} = 1$$

$$\sum_{x=1}^{\infty} \frac{3}{10^x} = \frac{3}{10} + \frac{3}{100} + \frac{3}{1000} + \dots$$

$$= 0.3 + 0.03 + 0.003 + \dots$$

$$= 0.33\bar{3} = \frac{1}{3}$$

Sum of a finite arithmetic sequence

- If $a_k = \{a_1, a_2, a_3, \dots, a_n\}$, then

$$\sum_{k=1}^n a_k = a_1 + a_2 + \dots + a_n$$

$$= n \left(\frac{a_1 + a_n}{2} \right)$$

$$\left(\frac{b_1 + b_2}{2} \right) h$$

Where: a_1 is the initial term

a_n is the final term


n is the number of terms

d is the common difference

Sum of a finite arithmetic sequence

- Example: A lecture hall has 8 seats in the first row and 24 in the last row. If each row has 2 more seats than the one in front of it, how many total seats are in the room?

$$a_1 = 8 \quad a_n = 24 \quad d = 2 \quad n = ????????$$



$$\text{find } n: a_n = a_1 + (n - 1)d$$

$$24 = 8 + (n - 1)(2)$$

$$24 = 8 + 2n - 2$$

$$24 = 2n + 6$$

$$18 = 2n$$

$$9 = n$$

$$\sum_{k=1}^n a_k = 9 \left(\frac{8 + 24}{2} \right)$$

$$= 9 \left(\frac{32}{2} \right)$$

$$= 9(16)$$

$$= 144$$

Sum of a finite geometric sequence

- If $a_k = \{a_1, a_2, a_3, \dots, a_n\}$, then

$$\begin{aligned}\sum_{k=1}^n a_k &= a_1 + a_2 + \dots + a_n \\ &= \frac{a_1(1-r^n)}{1-r}\end{aligned}$$

Where: a_1 is the initial term

n is the number of terms

r is the common ratio

Sum of a **finite** geometric sequence

- Example: Find the sum of: $4, -\frac{4}{3}, \frac{4}{9}, -\frac{4}{27}, \dots, 4\left(-\frac{1}{3}\right)^{10}$

a, r
 $a_1 = 4 \quad r = -\frac{1}{3} \quad n^{\text{th}} \text{ term is } 4\left(-\frac{1}{3}\right)^{10}, \text{ so } n = 11$

$$\begin{aligned} \sum_{k=1}^{11} a_k &= \frac{4\left(1 - \left(-\frac{1}{3}\right)^{11}\right)}{1 - \left(-\frac{1}{3}\right)} = \frac{4\left(1 - \left(-\frac{1}{3}\right)^{11}\right)}{\frac{4}{3}} = \frac{1 - \left(-\frac{1}{3}\right)^{11}}{\frac{1}{3}} \\ &= 3\left(1 - \left(-\frac{1}{3}\right)^{11}\right) \approx 3.000016935 \end{aligned}$$

Sum of an infinite geometric sequence

- An **infinite** geometric series **converges** if $|r| < 1$

If this is the case, then the sum of the series is:

$$\sum_{k=1}^{\infty} a_k r^{k-1} = \frac{a_1}{1-r}$$

Where: a_1 is the initial term

r is the common ratio

Sum of an infinite geometric sequence

- Find the sums (if the series converges):

1. $\sum_{k=1}^{\infty} 3(0.75)^{n-1}$ $|r|=0.75 < 1$, so it converges $a_1 = 3$, so

$$\sum_{k=1}^{\infty} 3(0.75)^{n-1} = \frac{a_1}{1-r} = \frac{3}{1-0.75} = \frac{3}{0.25} = 12$$

2. $\sum_{x=1}^{\infty} \frac{3}{10^x} = \frac{\frac{3}{10}}{1 - \frac{1}{10}} = \frac{\frac{3}{10}}{\frac{9}{10}} = \frac{3}{9} = \frac{1}{3}$

Homework 😊

- Pages 749-750;
 - Problems 1-21 odds, 25, 27, 29

